

NORTHERN ARIZONA PROVINCE (024)

By W.C. Butler

INTRODUCTION

This province covers about 59,000 sq mi, mostly in the southwestern part of the Colorado Plateau. Significant geologic features include the Grand Canyon, Kaibab Arch, Mogollon Highlands transition zone, Monument Uplift, Defiance Uplift, Black Mesa Basin, Holbrook Basin, and southern edges of the Kaiparowits and Blanding Basins.

The stratigraphic section shown for northeastern Arizona has demonstrated the highest petroleum potential in Arizona. See Wilson (1962), Butler (1988a), and Dickinson (1989) for synopses of the province's geology and evolution.

The lithologically and structurally complex basement of the Colorado Plateau area evolved from northwest-younging Proterozoic terranes sequentially accreted onto the Archean craton. As much as 12,000 ft of Middle and Late Proterozoic strata is preserved in possible rift-aulacogen depositional settings in central Arizona. Thick, unmetamorphosed, organic-rich Late Proterozoic strata deposited in backarc basins or continental lakes of north-central Arizona and south-central Utah have good petroleum potential. The plateau area, as a passive Paleozoic plate margin and buffered Mesozoic retro-arc platform, has been remarkably tectonically stable during Phanerozoic time.

The area is characterized by blanket Paleozoic strata, as much as 6,000 ft thick, consisting of mostly shallow marine clastics and carbonates showing numerous disconformities. These strata accumulated during transgressions and regressions from both the northwest and southeast, overlapping and thinning toward the trans-continental arch - a northeast-trending positive area extending from the northeast into central Arizona. Convergence between North and South American tectonic plates, with reactivation of basement blocks, during the late Paleozoic created the plateau's fault-bounded basins and uplifts. One of these downwarps, the Blanding Basin of the Four Corners area, is a prolific petroleum producer from the Pennsylvanian Hermosa Group. All Hermosa Group plays, including Arizona's largest accumulation--and one of the most notable fields in the United States, Dineh-bi-Keyah--are assessed in the Paradox Basin Province (021).

Paleozoic strata are overlain by complexly intercalated Mesozoic strata that generally become thinner from Black Mesa Basin toward the Mogollon Rim highlands, where

older strata crop out. Mesozoic strata were deposited in shallow marine shelf plus continental environments (coastal plain, piedmont, fluvial alluvial, eolian, lacustrine, and paludal). An apron of superjacent Oligocene to Pliocene volcanic flow rocks marks the southern province boundary; Pliocene and Quaternary volcanic fields (San Francisco Peaks) mark the center of the province. Incised rivers, such as the Colorado and Little Colorado Rivers, and their steep canyons are spectacular Quaternary features of the northern province. Flushing of hydrocarbons is a detrimental factor in these rapidly downcut areas. Late Tertiary erosion of the plateau with unloading of the crust, and consequential uplift, is another detriment because potential reservoirs are rendered underpressured.

Structurally, the Colorado Plateau is typified by long, northwest-trending Laramide monoclines and other gentle crustal warps; small fold axes are 10-15 mi long, but most are 30-65 mi long, and some even extend to 80 mi. A section of the Cordilleran hingeline may be present in the northwesternmost part of the province; platform and miogeosynclinal Paleozoic strata and Mesozoic continental to coastal strata comprise its stratigraphic section. The hingeline is considered the southern extension of the Sevier Thrust Belt of Utah; strata here were thrust eastward during the Late Cretaceous followed by regional extension during the Neogene and Quaternary. On the basis of seismic data, the basement depth in the resultant extensional basins is known to be 26,000 ft below sea level. Because this area has Great Basin physiography, it was not considered in the Colorado Plateau Province. Farther east, to the western edge of the Kaibab Arch, high-angle north-trending normal faults cut the plateau; they exhibit displacements of as much as several thousand feet. The net effect of this crustal extension is growth of the Great Basin at the expense of the Colorado Plateau and possible rupture of older traps.

The southern part of this province represents a structural transition from the saddles, troughs, benches and arches of the gentle northeast-dipping Mogollon Slope to an extremely complex horst-graben or half graben detachment fault regime. The zone exposes uplifted and dissected Precambrian igneous, metamorphic, and sedimentary rocks next to Tertiary basin deposits and has very low to no petroleum potential. During the Jurassic and Cretaceous Periods this transition zone became the edge of a magmatic arc fully developed in southwestern Arizona (active subduction to the southwest). The arc was a highland source area; streams drained northeastward depositing up to 5,000-7,500 ft of sediment in northern Arizona.

A cluster of 10 small oil and gas fields, discovered between 1954 and 1971, produce from structural traps in Devonian, Mississippian, and Pennsylvanian strata in the extreme northeastern part of Arizona. The three largest of these fields (Dry Mesa, East Boundary Butte, and Teec Nos Pos) have a combined cumulative production of about 2.2 MMBO and condensate, and 12 BCF of gas. Twenty-five mi south of these fields is Dineh-bi-Keyah (1967) which is the province's only oil field of significant size; it has an estimated ultimate recovery of 20 MMBO from Oligocene syenite sills. Sill reservoirs were very locally sourced by Pennsylvanian black shale and dolomite matured by the heat of the igneous intrusions. Cumulative production through 1993 for all fields in Province 024 is 20 MMBO and condensate; about 25 BCF of flammable gas; and, 9.3 BCF of helium. Methodical exploration of the province is inadequate; most wells were drilled as promotional ventures rather than on the concepts of modern petroleum geology. Some large structures have never been drilled, and the deeper reservoirs are rarely tested. Depths in feet for the deepest boreholes in each of the seven northernmost counties (including northern Apache County, which is the only producing area to date) are 0, 480 ft, 3,206 ft, 7,070 ft, 7,210 ft, 7,780 ft, and 8,461 ft. Drilling density is about 155 sq mi per borehole, including Apache County, and about 420 sq mi per borehole, excluding the productive area.

Four plays were assessed. The Hurricane Fault-Uinkaret Plateau Play (2401) was assessed qualitatively only; the other three plays were assessed quantitatively and include the Late Proterozoic (Chuar-Sourced) and Lower Paleozoic Play (2403), the Oraibi Trough Play (2402), and the Holbrook Basin Anticline and Stratigraphic Traps Play (2404).

ACKNOWLEDGMENTS

Scientists affiliated with the American Association of Petroleum Geologists and from various State geological surveys contributed significantly to play concepts and definitions. Their contributions are gratefully acknowledged.

CONVENTIONAL PLAYS

2401. HURRICANE FAULT-UINKARET PLATEAU PLAY (HYPOTHETICAL)

This oil and associated-dissolved gas play of northwestern Arizona is unproductive, hypothetical, and highly speculative. The play is in north-central Mohave County north of the Colorado River and almost entirely east of the Hurricane Fault. Drag and reverse drag folds along the Hurricane Fault and gentle plateau folds provide structural traps. The play is based on evidence of hydrocarbons in the system, such as numerous live shows in boreholes of the region (particularly from Pennsylvanian-Permian rocks), plus common strong fetid odors, stains, seeps, and tar in outcrops. The abandoned shallow Virgin oil field 15 mi north produced minor amounts of oil from the Triassic Moenkopi Formation.

Reservoirs: Potential reservoirs are Devonian (Martin and Temple Butte Formations), Mississippian (Redwall Limestone), Pennsylvanian (Callville Limestone and Supai Groups), and Permian (Supai Group, Queantoweap and Esplande Sandstones, Coconino Sandstone, and Toroweap Formation). These formations have all produced good hydrocarbon shows. The Temple Butte is an olive-gray, silty and sandy supratidal, irregularly-bedded carbonate. The Redwall is a massive, light-gray, shallow, open-marine, equatorial carbonate. The 1,000-ft-thick Supai is a complex set of mostly clastic beds that are light-gray to red, calcareous and gypsiferous mudstones, siltstones, and cross-bedded sandstones of estuarine to intertidal tropical origin. The Hermit consists of reddish fine clastics deposited in distal fan, floodplain, and fluvial environments. The Coconino is an eolian, well-sorted, cross-bedded and rippled, buff quartz sandstone. The Toroweap has intercalated fine-grained, contorted sandstones, siltstone, mudstones, dolomite, gypsum, and fossiliferous cherty limestone representing deposition in a low-energy marine to sabkha environment. As much as 2,000 to 2,500 ft of this total stratigraphic section may have reservoir potential.

Source rocks: Source rocks probably containing type I organic matter now in the thermal zone of oil generation may include the mudstones and carbonates of the same above-noted formations, but also include the Kaibab Limestone (gray, fossiliferous, cherty, aphanitic) and, less likely, the Hermit Shale; however, stronger evidence for better source rocks is lacking and hence degrades the petroleum potential.

Traps: Recurrent Miocene and younger normal faulting of a Laramide-formed monocline formed drag and reverse-drag folds along the high-angle Hurricane Fault (east side up and west side down). Southwestward updip migration of hydrocarbons

may have been trapped in these folds or against the fault. Gentle northwest-trending anticlines to the east on the Uinkaret Plateau (Giardina, 1979) have not been adequately explored, and possible tilted oil-water contacts may have prevented discovery of accumulations in those folds that have been drilled. Targets are most likely 3,500-4,500 ft deep, but conceivably could cover a wide range of from 500 to 6,000 ft deep. Migration time is unknown but was probably Laramide.

Numerous unconformities, porosity-permeability pinchouts, and fine clastic units (for example, Hermit Shale and Moenkopi Formation) are potential seals. Gypsiferous beds of Permian age could provide additional seals.

Exploration status and resource potential: Methodical exploration has been lacking. Less than 25,000 ft of strata have been drilled in all boreholes of this play. Borehole control is sparse and irregularly distributed, averaging about one per 400 sq mi. Some major folds have been tested but none adequately. The ratio of good shows to dry holes is greater than 1.0 for boreholes 1,000-5,000 ft deep, and hence the area is encouraging as a result of the shows, good reservoir qualities, and abundant structural and stratigraphic traps

2402. ORAIBI TROUGH PLAY

This confirmed oil and associated gas play is mostly in northeastern Arizona but extends into southeastern Utah, southwestern Colorado, and northwestern New Mexico between the Defiance Uplift on the east and the Monument Uplift and Kaibab positive areas on the west. Although it is a confirmed play, 99 percent of the play is south of the production in Devonian, Mississippian, and Pennsylvanian strata of northeasternmost Arizona. The basis of the play is a close association of Late Devonian and Early Mississippian source and reservoir strata with northwest-trending anticlines and stratigraphic traps in the northeast-trending Oraibi Trough, which formed in mid-Paleozoic time. This play does not consider Hermosa Group production from Pennsylvanian carbonate-mound reservoirs on the Arizona part of the southern shelf of the Blanding-Paradox Basin, which is assessed in province 021.

Geographically, the center of the Oraibi Trough nearly coincides with the greater Black Mesa structural basin formed during Laramide time. The total stratigraphic southeast-thinning section from outcrop to basement is 4,000-9,000 ft thick roughly, 40 percent is Paleozoic foreland restricted-shelf strata, and the remainder is a mix of continental to shallow-marine strata of Mesozoic age. The maximum combined thickness of the

Devonian-Mississippian section is about 1,000-1,250 ft (Beus, 1989). Pennsylvanian strata (as much as 1,800 ft thick) are a secondary consideration for reservoir and source rocks in this play. Strata of all three periods pinch-out against the Defiance Uplift.

Reservoirs: The best reservoirs are in relatively small but widespread deltaic clastic wedges or lenses of the Upper Devonian McCracken Sandstone Member (as much as 250 ft thick) of the Elbert Formation (Kashfi, 1983). These medium- to coarse-grained and medium- to well-sorted sand bodies splay westward off the Defiance erosional basement into the trough and are encased by less porous and less permeable strata deposited during east-west transgressions and regressions. The McCracken Sandstone and the pelletal dolomite, skeletal and pelletal carbonate mudstone, and sandy dolomite of the upper part of the Elbert Formation are productive in the Paradox Basin. Also having reservoir potential are the superjacent porous, cherty, crinoidal and pelletal Mississippian carbonates and the Pennsylvanian Hermosa Formation and Supai Group clastics. Seals include waxy shale and anhydrite of the Elbert Formation, or dense massive limestones and dolomites of the Redwall Formation and Leadville Limestone, or redbeds, calcareous siltstones and mudstones, and thin gypsum units of the Triassic Moenkopi Formation.

Source rocks: The best source rock may be the Aneth Formation, which is about 85 percent within the oil-generation window having maturities of R_o 0.90-2.1 percent ; it probably contains 0.50-1.03 percent TOC of type II and III (Tucker, 1983) in the richest areas. Lithologically, it is a dark, glauconitic, argillaceous aphanitic dolomite interbedded with dark carbonaceous shale and siltstone and minor limestone and evaporites. Given outcrops having thermal maturities of R_o 0.41-0.60 percent, the floor of the oil window is in Devonian rocks and may range from +3,000 ft to -500 ft in elevation. Shallower Pennsylvanian strata may have been matured by igneous intrusions of Tertiary age, such as at the Dineh-bi-Keyah oil field on the Defiance Uplift. Geophysical signatures (gravity and magnetic), suggesting shallow intrusions, similar to the Dineh-bi-Keyah area, can be found along the perimeter of the Defiance, that is, the eastern margin of this play. Intrusive centers, such as Hopi Buttes and the Carrizo Mountains, are excellent areas to evaluate for structural traps associated with source rock matured by the heat of cooling igneous rock. In addition to the significant live shows in boreholes, the Devonian rocks in this play, and particularly in outcrops along the Mogollon Slope, are known for their strong fetid odor, seeps, asphalt-filled fractures, and other hydrocarbon impregnations.

Traps: Both structural and stratigraphic traps are plentiful. Total structural relief in the play is 6,000-7,000 ft, providing good flow gradients up the Mogollon Slope to the south and towards the east-west margins of the trough, particularly toward the Defiance Uplift. Subtle long-distance facies changes and fluid migration probably characterize northeastern Arizona. Up-dip facies changes and stratigraphic wedge-outs are prevalent. Vuggy biohermal or other carbonate mounds may present yet another highly probable trap in the Mississippian and Pennsylvanian strata. Significant unconformities between Mississippian and Pennsylvanian strata and between Pennsylvanian and Triassic strata also provide a speculative trapping mechanism. The cumulative axial lengths of major anticlines is 350 mi and that of monoclines is 225 mi. Structural closure on the upfolds is 500-1,000 ft. Regional Jurassic tilting of the Colorado Plateau in this area may have displaced the oil-water contacts off the structural center. Drilling depth to all traps is 3,300-7,500 ft.

Timing and generation: Timing of generation and migration is not known, but could have reasonably occurred from early Cretaceous to early Tertiary time; hence, migration could have been prior to structural trap formation. Present-day heat flow is 1.13-1.80 HFU, and the present-day geothermal gradient is 1.05-1.90 F/100 ft.

Exploration status: Because about 98 percent of the play is on Native American land, exploration essentially did not begin until the mid-1960's, when it was first permitted. Cumulative production from Devonian and Mississippian reservoirs has been 908,000 BO and 1.3 BCF of gas. The Redwall Limestone at Dry Mesa field has produced about 802,000 BO from fractured dolomite on an anticline that has 35 ft of closure. Average borehole depth in the play is 4,350 ft. Drilling density in the play is about one borehole per 185 sq mi; less than 2 percent of the area has been drilled to the Coconino Sandstone. Less than 1 percent of the area has been drilled to the Devonian. Drilling density is much higher, however, if only the northeasternmost part of the play is considered, where density is one borehole per 6-7 sq mi to an average depth of 5,770 ft. The play is grossly under-explored, and many obvious large upfolds are untested.

Resource potential: The probability that this play "works" (a product of the all attributes) is about 40 to 45 percent for oil and associated gas. The median size of undiscovered accumulations is 1.5 MMBO, and the probable median number of accumulations is 5.

2403. LATE PROTEROZOIC (CHUAR-SOURCED) AND LOWER PALEOZOIC PLAY (HYPOTHETICAL)

About one-third of the area of this unusual hypothetical oil and thermal gas play is in north-central Arizona and two-thirds in south-central Utah of the Paradox Basin Province 021 (see play 2108). The play is hypothetical and highly speculative, being defined on the inference that the Late Proterozoic Chuar Group sourced reservoir units within itself and in superjacent Paleozoic reservoirs, primarily the Cambrian Tapeats Sandstone. Measured thickness of the group in the eastern Grand Canyon ranges from 5,370 to 6,400 ft. Potential accumulations are associated with large Colorado Plateau structural traps (see province description). Boundaries for this play are ill-defined because so few boreholes have penetrated the Chuar; consequently, its regional occurrence and facies are poorly understood. See Rauzi (1990) for additional information.

Reservoirs: Reservoirs, probably underpressured, include numerous fine clastic (mostly siltstone) beds in the Chuar Group (if sufficiently fractured), the superjacent sandstone of the Sixtymile Formation, the Tapeats Sandstone (150 to 400-ft-thick cross-bedded beach deposit), and possibly other Paleozoic sandstones if extensive vertical migration has occurred. There have been significant petroleum shows in every Paleozoic system and also in Triassic strata. Multi-billion-barrel-size accumulations of hydrocarbons are present in the Permian and Triassic "tar sand triangle" and the Circle Cliffs-White Canyon areas of south-central Utah. Their origin is a long-standing enigma; some explorationists suggest the Chuar Group may have been their source, but this hypothesis remains as only one of several possible explanations. There is one field in this play, the Upper Valley oil field (discovered 1964) of south-central Utah that has produced more than 20 MMBO from Permian and Triassic reservoirs.

Source rocks: The Walcott Member (Elston, 1989) is organic-rich gray to black mudstone and siltstone containing thin sequences of sandstone and stromatolitic crypto-algal, pisolitic dolomite probably deposited in lacustrine to tidal flat environments. Stratigraphically, where measured, the 850- to 1,100-ft-thick Walcott Member of the Chuar Group is the youngest member of the Kwagunt Formation and is, according to current knowledge, the richest unit. Total organic carbon (algal type) has been documented as high as 10 percent in the dark mudstones; 3 percent TOC is average for the source-rock component of the Walcott. Thermal maturity of the Chuar Group is partially within the assumed oil-generation window, (R_o 0.8-1.35 percent). Perhaps half of the mature source rock is greater than R_o 1.35 percent (the oil floor) and

in the gas generative phase. Upper Paleozoic and Mesozoic outcrops in the play area have R_o values between 0.5 and 0.6 percent, indicating that they have not been deeply buried. In the Arizona part of the play, the top of the Chuar Group is probably at R_o 0.8-1.0 percent.

Drilling depths to potential accumulations are from 2,000 to 13,000 ft (average 6,500 ft) for oil and from 6,000 to 20,000 ft (average 9,000 ft) for gas. Depth to the top of the Chuar Group at the Utah-Arizona state line in the center of the play is about 13,000-13,500, ft implying that Arizona is oil-prone and Utah is gas-prone. The Utah source beds are more mature, having been buried deeper in the Kaiparowits Basin area than in the Arizona Kaibab Plateau area. Chuar depths on the west side of the play near the Arizona-Utah state line are relatively shallow, at about 5,000 ft, due to the north-trending Kaibab uplift.

Timing and migration and Traps: Timing of generation and migration are two big unknowns in this play. Laramide is suspected as being reasonable for timing, but questions remain. If migration occurred prior to the Late Paleozoic uplifts or barriers, accumulations may be a long distance from the play area. If migration occurred after Late Paleozoic and Laramide deformations, ample structural traps would be available. Stratigraphic traps (porosity and permeability pinchouts and major unconformities and angular unconformities) are also known to be present but are not necessary to make this play work. Within this play the combined axial lengths of anticlines is 552 mi, and that of monoclines is 464 mi. Drilling of large anticlines has been unsuccessful to date; many large anticlines are undrilled; and the wells that have been drilled may have missed accumulations due to unusual hydrologic situations (Allin, 1990). These structures are large enough to permit very large accumulations. Graben-like structures in the basement aulocogen (?) could also trap hydrocarbons. The Bright Angel Shale is probably an excellent seal overlying the Tapeats. Other seals in the Paleozoic section include impermeable carbonates, fine clastics, and evaporites.

Exploration status and resource potential: If the Chuar is a good source rock as currently believed, one would expect to find relic oil or seeps in this play; however no Chuar seeps have been reported in the Grand Canyon or in the paleo high areas, such as around the Kaibab Arch. On the other hand, more than a dozen shows have been encountered in the Paleozoic section in Arizona west of longitude 112°. Some of the Cambrian penetrations in the play encountered only the Bright Angel Shale and Mauv Limestone but did not test the lower potential reservoir in the Tapeats Sandstone; thus

exploration to the Chuar is nearly nonexistent. The Tidewater Kaibab Federal 1-A well (logged 1957), Kane County, Utah, found oil and gas shows in both the upper Tapeats Sandstone (15 ft of oil at a depth of about 4,850 ft) and in the Chuar Group (at a depth of about 5,900 ft). Other Chuar tests are currently being permitted or drilled.

2404. HOLBROOK BASIN ANTICLINE AND STRATIGRAPHIC TRAPS PLAY (HYPOTHETICAL)

The Holbrook Basin is a nonproducing hypothetical oil and associated gas play in east-central Arizona. The Little Colorado Rivers flow northwest through the center of the basin. The play has a structural component (the northwest-trending Holbrook Anticline) near the southwest basin margin, and a stratigraphic trapping component to the northeast. Rapid facies changes and updip pinchouts characterize the basin.

The shallow Holbrook Basin, in a shallow shelf environment, formed over a Precambrian depression beginning in the Pennsylvanian period, was filled with Permian sabkha sediments, and was later modified by mild Laramide tectonism. Maximum present-day structural relief is only 1,500 ft. About 400 ft of Devonian and Mississippian strata disconformably overlie the basement and are subjacent to the play. The thickest strata (3,000-4,250 ft) in the basin are cyclic, Pennsylvanian and Permian, interbedded clastics (including redbeds) and carbonates (brown dolomites and dark micritic limestones) with basin-centered evaporites (anhydrite, halite and potash) and lesser dark, calcareous shales; these strata comprise both the reservoir and source rocks. Specifically, the Horquilla Limestone and Naco Formation (900 ft thick), Supai Group (900-1,350 ft thick), and the Coconino, Glorieta, and DeChelly Sandstones (1,000-2,000 ft) are potential reservoirs. About 70-85 percent of the Pennsylvanian section and 90-95 percent of the Permian section is clastic. These units have good porosity and are disconformably overlain by Triassic outcrops; Quaternary basalt, which had no adverse affect on the source rocks, covers the extreme southeastern part of the basin. Production of helium (probable Precambrian source) from a faulted anticline at nearby Pinta Dome indicates that the Permian reservoirs have very little leakage. The thick salt mass of the basin is used to store commercial natural gas (Peirce, 1981), and demonstrates effective seals.

Source rocks, timing, and migration: Source rocks are primarily the Naco Group and Supai Group, which includes the thin, black Fort Apache Limestone. The evaporite facies of the Supai Group may have preserved much of its mixed type I and II organic matter. Regionally, limited testing has shown TOC values in the carbonates of the Supai to be at least 1-2 percent. In late 1993, 40 mi southeast of the Holbrook Basin the

Alpine-Federal #1 geothermal test well encountered an excellent live (bleeding) oil show and two other significant indications of hydrocarbons in the Supai Group (Rauzi, 1994). The intrusive igneous dikes found in this section had virtually no thermal effect on the host strata. The entire Late Paleozoic section is within the oil-generation window and ranges from R_o 0.65-1.1 percent. Present-day geothermal gradients range from 1.2 to 1.6; F/100 ft. Present-day heat flux immediately surrounding the basin is much higher; these areas have good geothermal resource potential. Timing for generation and migration is unknown, but it was probably during the Laramide.

Traps: Depth to traps is 500-4,000 ft; tops to evaporites within the Supai Group occur at depths of 1,000-2,450 ft. Traps include the Holbrook Anticline, which has an axial length of about 60 mi and a maximum closure of about 250 ft. The anticline may be a result of either evaporite dissolution (collapse structure) or mild Laramide northeast-southwest compression, or both. Other secondary folds trend northeast and may also trap hydrocarbons. Northeast updip migration of petroleum could also be trapped stratigraphically. The Paleozoic units thin and pinch-out toward the Defiance positive area; the magnitude of the disconformities increases in that direction providing various mechanisms for accumulation. Interfingering evaporite or impermeable carbonate beds could seal these accumulations.

Exploration status and resource potential: Although drilling in the Holbrook Basin has been frustrating, it is also promising because of the high number of live oil and gas shows and strong fetid odors in Late Paleozoic outcrops of the area. Abundant shows occur at depths between 1,100-3,200 ft. Some boreholes have encountered as many as eight shows. Oil stains, asphaltic oil, and dead heavy oil have been encountered in many wells. However, even with these shows, proof of an adequate source rock, and the possibility of fresh-water flushing, inhibit a higher exploration effort. The crest of the Holbrook Anticline may be offset at depth, and the more recent off-surface-structure boreholes seem to find the best shows. The basin has a drilling density of about one borehole per 45 sq mi. Many of the wells were drilled specifically for potash. Probably less than 20 boreholes have been drilled below 3,600 ft. Many explorationists in Arizona have observed that wells tend to be drilled haphazardly, and the drilling in the basin has been inconclusive with many structures left to be tested. Although this is a high-risk frontier area, leasing and drilling in the last 10 years have been fairly encouraging.

UNCONVENTIONAL PLAYS

There are no unconventional plays described in this province report. However, unconventional plays listed in the surrounding provinces may include parts of this province. Individual unconventional plays are usually discussed under the province in which the play is principally located.

REFERENCES

- Aiken, C.L.V., Laughlin, A.W., and West, F.G., 1978, Residual Bouguer gravity anomaly map of New Mexico: University of California Los Alamos Scientific Laboratory LA-7466-map/UC-11, scale 1:500,000.
- Aiken, C.L.V., and Sumner, J.S., 1974, A geophysical and geological investigation of potentially favorable areas for petroleum exploration in southeastern Arizona: Arizona Oil and Gas Conservation Commission Report of Investigations 3, 40 p.
- Allin, D.L., 1990, Colorado Plateau subsurface water flow key: Oil and Gas Journal, v. 88, no. 30, July 23, p. 52-54.
- American Geophysical Union, 1989, Sedimentation and tectonics of western North America, v. 3: Washington, D.C., American Geophysical Union, 28th International Geological Congress, 477 p.
- Anderson, O.J., Lucas, S.G., Love, D.W., and Cather, S.M., eds., 1989, Southeastern Colorado Plateau: New Mexico Geological Society, 40th Field Conference Guidebook, 348 p.
- Anderson, R.Y., and Harshbarger, J.W., eds., 1958, Black Mesa Basin--Guide book of northeastern Arizona: Arizona Geological Society, 9th Field Conference, 205 p.
- Anschutz, P.F., 1980, The overthrust belt--Will it double U.S. gas reserves?: World Oil, v. 190, no. 1, p. 111-116.
- Arizona Oil and Gas Conservation Commission, 1961, Oil, gas and helium in Arizona: Phoenix, Arizona, Arizona Development Board, 108 p.
- Arizona Oil and Gas Conservation Commission, 1987, State of Arizona well location map four: Phoenix, Arizona, August, 38 p.
- Arizona Oil and Gas Conservation Commission, 1992, Arizona well location map and report: Arizona Geological Survey Oil and Gas Publication OG-12, 22 p., scale 1:500,000.
- Armentrout, J.M., Cole, M.R., and TerBest, Harry, Jr., eds., 1979, Cenozoic paleogeography of the western United States: Society of Economic Paleontologists and Mineralogists, Pacific Coast Section Symposium 3, 335 p.
- Armstrong, A.K., Kottowski, F.E., Stewart, W.J., Mamet, B.L., Baltz, E.H., Siemers, W.T., and Thompson, Sam, III, 1979, The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States--New Mexico: U.S. Geological Survey Professional Paper 1110-W, 27 p.

- Arnold, E.C., and Hill, J.M., comps., 1981, New Mexico's energy resources '80: New Mexico Bureau of Mines and Mineral Resources Circular 181, 59 p.
- Baars, D.L., ed., 1979, Permianland: Four Corners Geological Society Guidebook, 9th Field Conference and Field Symposium, 186 p.
- Baars, D.L., 1983, The Colorado Plateau--A geologic history: Albuquerque, New Mexico, University of New Mexico Press, 279 p.
- Bahr, C.W., 1962, The Holbrook anticline, Navajo County, Arizona, *in* Weber, R.H., and Peirce, H.W., eds., Guidebook of the Mogollon Rim region east-central Arizona: New Mexico Geological Society, 13th Field Conference, p. 118-122.
- Barwin, J.R., 1969, Oil possibilities in Arizona: Oil and Gas Journal, v. 67, no. 35, p. 186-188.
- Barwin, J.R., King, R.W., and Hassenfratz, C.A., 1971, Future oil and gas potential of northeast Arizona, *in* Cram, I.H., ed., Future petroleum provinces of the United States--Their geology and potential: American Association of Petroleum Geologists Memoir 15, p. 449-469.
- Bayer, K.C., comp., 1983, Generalized structural, lithologic and physiographic provinces in the fold and thrust belts of the United States: U.S. Geological Survey map, scale 1:2,500,000.
- Bayley, R.W., and Muehlberger, W.R., 1968, Basement rock map of the United States: U.S. Geological Survey Map, scale 1:2,500,000.
- Bayliss, G.S., and Schwarzer, R.R., 1992, New Mexico hydrocarbon source rock evaluation project--New Well Peak section in Big Hatchet Mountains, Hidalgo County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 379, (revised January, 1992), unpaginated.
- Beal, C.H., 1948, reconnaissance of the geology and oil possibilities of Baja California, Mexico: Geological Society of America Memoir 31, 138 p.
- Beatty, Barbara, and Wilkinson, P.A.K., eds., 1986, Frontiers in geology and ore deposits of Arizona and the Southwest: Arizona Geological Society Digest, V. XVI, 554 p.
- Beikman, H.M., Peterson, J.A., Huber, D.F., and Butler, W.C., 1986, Metallic mineral and mineral-fuel resource potential map of Arizona showing major mineral deposits: U.S. Geological Survey Map Mineral Investigations Resource Map MR-94, scale 1:1,000,000.

- Beus, S.S., 1989, Devonian and Mississippian geology of Arizona, *in* Jenney, J.P., and Reynolds, S.J., eds., *Geologic evolution of Arizona: Arizona Geological Society Digest, V. 17*, p. 287-311.
- Billingsley, G.H., 1978, A synopsis of stratigraphy in the western Grand Canyon: Flagstaff, Arizona, Museum of Northern Arizona Research Paper 16, 26 p.
- Bilodeau, W.L., and Lindberg, F.A., 1983, Early Cretaceous tectonics and sedimentation in southern Arizona, southwestern New Mexico and northern Sonora, Mexico, *in* Reynolds, M.W., and Dolly, E.D., eds., *Mesozoic paleogeography of west-central United States: Society of Economic Mineralogists and Paleontologists, Symposium 2*, p. 173-188.
- Blackwell, D.D., 1978, Heat flow and energy loss in the western United States, *in* Smith, R.B., and Eaton, G.P., eds., *Cenozoic tectonics and regional geophysics of the Western Cordillera: Geological Society of America Memoir 152*, p. 175-208.
- Blakey, R.C., 1979, Oil impregnated carbonate rocks of the Timpweap Member Moenkopi Formation, Hurricane Cliffs area, Utah and Arizona: *Utah Geology*, v. 6, p. 45-53.
- Brady, B.T., 1984, Selected geologic and hydrologic characteristics of the Basin and Range Province, western United States, coal, oil and gas wells, seeps and tar sandstone occurrences: U.S. Geological Survey Miscellaneous Investigations Series Map I-1522-E, scale 1:2,500,000.
- Branson, C.C., ed., 1962, Pennsylvanian system in the United States--A symposium: American Association of Petroleum Geologists, 508 p.
- Brennan, D.J., 1989, Hydrocarbon possibilities in Yuma area, Arizona [abs.]: *American Association of Petroleum Geologists Bulletin*, v. 73, no. 3, p. 337.
- Broadhead, R.F., and Black, B.A., 1989, Petroleum exploration wells drilled in west-central New Mexico, *in* Anderson, O.J., and others, eds., *South eastern Colorado Plateau: New Mexico Geological Society, 40th Field Conference Guidebook*, p. 287-296.
- Bryant, D.L., 1968, Diagnostic characteristics of the Paleozoic formations of southeastern Arizona, *in* Titley, S.R., ed., *Southern Arizona guidebook III: Tucson, Arizona, Arizona Geological Society*, p. 33-47.
- Butler, W.C., 1971, Permian sedimentary environments in southeastern Arizona, *in* McCullough, E.J., ed., *Arizona Geological Society Digest, V. IX*, p. 71-94.

- Butler, W.C., 1985, Petroleum potential of the Hualapai Indian Reservation in northwestern Arizona--A preliminary qualitative assessment for the tribe: unpublished U.S. Geological Survey and Bureau of Indian Affairs Administrative Report, on file at Bureau of Indian Affairs, 20 p.
- Butler, W.C., 1988a, The rationale for assessment of undiscovered, economically recoverable oil and gas in central and northern Arizona Play analyses of seven favorable areas: U.S. Geological Survey Open-File Report 87-450-V, 145 p.
- Butler, W.C., 1988b, The rationale for assessment of undiscovered, economically recoverable oil and gas in south-central New Mexico--A geologic overview and play analysis of two favorable areas: U.S. Geological Survey Open-File Report 88-450-B, 134 p.
- Butler, W.C., 1989a, The geologic setting of southern Arizona and southwestern New Mexico, with a rationale for assessment of undiscovered, economically recoverable oil and gas--A summary of four potential plays: U.S. Geological Survey Open-File Report 88-450-M, 150 p.
- Butler, W.C., 1989b, Strategic crude oil storage in Arizona evaporite masses--An initial description, evaluation, and selected references of four potential sites: unpublished U.S. Geological Survey Administrative Report, on file with the Branch of Petroleum Geology, Denver, Colorado, 109 p.
- Butler, W.C., 1992, Elevation data for the Precambrian surface in the central and southern Colorado Plateau and vicinity: U.S. Geological Survey Open File Report 92-194, 69 p.
- Callender, J.F., Wilt, J.C., and Clemons, R.E., eds., 1978, Land of Cochise Southeastern Arizona: New Mexico Geological Society Guidebook, 29th Field Conference, 372 p.
- Cather, S.M., and Johnson, B.D., 1984, Eocene tectonics and depositional setting of west-central New Mexico and eastern Arizona: New Mexico Bureau of Mines and Mineral Resources Circular 192, 33 p.
- Chapin, C.E., ed., 1983, Socorro region II: New Mexico Geological Society, 34th Field Conference Guidebook, 344 p.
- Christiansen, R.L., and Lipman, P.W., 1972, Cenozoic volcanism and plate tectonic evolution of the western United States, II. Late Cenozoic: Philosophical Transactions of the Royal Society of London, Series A, v. 271, no. 1213, p. 249-284.
- Clemons, R.E., King, W.E., and Mack, G.H., eds., 1986, Truth or Consequences region: New Mexico Geological Society, 37th Field Conference Guidebook.

- Clemons, R.E., and Lawton, T.F., 1991, Petrographic and well log analysis of four exploration wells in southwestern New Mexico: New Mexico Bureau of Mines and Mineral Technology Open-File Report 381, 97 p.
- Condie, K.C., 1981, Precambrian rocks of the southwestern United States and adjacent areas of Mexico: New Mexico Bureau of Mines and Mineral Resources, Resource Map 13, scale 1:500,000.
- Condie, K.C., 1982, Plate-tectonics model for Proterozoic continental accretion in the Southwestern United States: *Geology*, v. 10, p. 37-42.
- Coney, P.J., 1979, Tertiary evolution of Cordilleran metamorphic core complexes, *in* Armentrout, J.M., Cole, M.R., and TerBest, H., Jr., eds., *Cenozoic paleogeography of the western United States*: Society of Economic Paleontologists and Mineralogists, Symposium 3, p. 15-28.
- Conley, J.N., and Giardina, S., Jr., 1979, Favorable and potentially favorable areas for hydrocarbon and geothermal energy sources in northeastern Arizona: Arizona Oil and Gas Conservation Commission, Report of Investigations 7, 56 p.
- Corbitt, L.L., and Woodward, L.A., 1973, Tectonic framework of Cordilleran foldbelt in southwestern New Mexico: *American Association of Petroleum Geologists Bulletin*, v. 57, p. 2207-2216.
- Cordoba, D.A., Wengerd, S.A., and Shomaker, John, eds., 1969, *Guidebook of the Border Region*: New Mexico Geological Society, 20th Field Conference, 218 p.
- Crysdale, B.L., and Schenk, C.J., 1988, Bitumen-bearing deposits of the United States: *U.S. Geological Survey Bulletin* 1874, 45 p.
- Dane, C.H., and Bachman, G.O., 1965, Geologic map of New Mexico: U.S. Geological Survey and New Mexico Bureau of Mines and Mineral Resources, scale 1:500,000.
- Davis, G.H., 1978, Monocline fold pattern of the Colorado Plateau, *in* Matthews, V., III, ed., *Laramide folding associated with basement block faulting in the western United States*: Geological Society of America Memoir 151, p. 215-233.
- Davis, G.H., 1979, Laramide folding and faulting in southeastern Arizona: *American Journal of Science*, v. 279, p. 543-569.
- Davis, G.H., and VandenDolder, E.M., eds., 1987, Geologic diversity of Arizona and its margins--Excursions to choice areas: Arizona Bureau of Geology and Mineral Technology, Geological Survey Branch Special Paper 5, 422 p.

- DeFord, R.K., and Kehle, R.O., Chairmen, 1976, Geothermal gradient map of North America: American Association of Petroleum Geologists and U.S. Geological Survey, scale 1:5,000,000.
- Desborough, G.A., Poole, F.G., Daws, T.A., and Scarborough, R.B., 1984, Hydrocarbon source rock evaluation of the Middle Proterozoic Apache Group Gila County Arizona, *in* Woodward, J., and others, eds., Hydrocarbon source rocks of the Greater Rocky Mountain region: Rocky Mountain Association of Geologists Symposium, p. 51-55.
- Dickinson, W.R., 1981, Plate tectonic evolution of the southern Cordillera, *in* Dickinson, W.R., and Payne, W.D., eds., Relations of tectonics to ore deposits in the southern Cordillera: Arizona Geological Society Digest, V. XIV, p. 113-135.
- Dickinson, W.R., 1989, Tectonic setting of Arizona through geologic time, *in* Jenney, J.P., and Reynolds, S.J., eds., Geologic evolution of Arizona: Arizona Geological Society Digest, V. 17, p. 1-16.
- Dickinson, W.R., and Klute, M.A., eds., 1987, Mesozoic rocks of southern Arizona and adjacent areas: Arizona Geological Society Digest, V. 18, 394 p.
- Dixon, G.H., Baltz, D.H., and Stipp, T.F., 1954, Records of wells drilled for oil and gas in New Mexico: U.S. Geological Survey Circular 33, 79 p.
- Drewes, H.D., 1981, Tectonics of southeastern Arizona: U.S. Geological Survey Professional Paper 1144, 96 p.
- Eberly, L.D., and Stanley, T.B., Jr., 1978, Cenozoic stratigraphy and geologic history of southwestern Arizona: Geological Society of America Bulletin, v. 89, p. 921-940.
- Elston, D.P., 1979, Late Precambrian Sixtymile Formation and orogeny at top of the Grand Canyon Supergroup, northern Arizona: U.S. Geological Survey Professional Paper 1092, 20 p.
- Elston, D.P., 1989, Grand Canyon Supergroup, northern Arizona--Stratigraphic summary and preliminary paleomagnetic correlations with parts of other North American Proterozoic successions, *in* Jenney, J.P., and Reynolds, S.J., eds., Geologic evolution of Arizona: Arizona Geological Society Digest, V. 17, p. 259-272.
- Fassett, J.E., ed., 1978, Oil and gas fields of the Four Corners area, v. I and II: Four Corners Geological Society, 727 p.
- Foster, R.W., 1964, Stratigraphy and petroleum possibilities of Catron County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Bulletin 85, 55 p.

- Foster, R.W., 1978, Oil and gas evaluation of White Sands Missile Range and Fort Bliss Military Reservation, south-central New Mexico: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 92, 60 p.
- Fouch, T.D., and Magathan, E.R., eds., 1980, Paleozoic paleogeography of the west-central United States: Society of Economic Paleontologists and Mineralogists, Symposium 1, 431 p.
- Frazier, W.J., and Schwimmer, D.R., 1987, Regional stratigraphy of North America: New York, New York, Plenum Press, 719 p.
- Fritz, M., 1990, Precambrian source sparks imaginations--Chuar inspires quest in the West: American Association of Petroleum Geologists Explorer, October, p. 8-9.
- Frost, E.G., and Martin, D.L., eds., 1982, Mesozoic-Cenozoic tectonic evolution of the Colorado River Region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, Inc., Anderson-Hamilton Volume, (published in conjunction with a symposium and field trip for the Geological Society of America), 608 p.
- Gehrels, G.E., and Spencer, J.E., eds., 1990, Geologic excursions through the Sonoran Desert region, Arizona and Sonora: Arizona Geological Society Special Paper 7, Field-Trip Guidebook, Cordilleran Section, Geological Society of America, 202 p.
- Gerrard, T.A., 1966, Environmental studies of Fort Apache Member, Supai Formation, east-central Arizona: American Association of Petroleum Geologists Bulletin, v. 50, p. 2434-2463.
- Giardina, S., Jr., 1979, Geologic review of northwestern Arizona for petroleum exploration investigators: Arizona Oil and Gas Conservation Commission, Special Publication No. 4, 72 p.
- Grant, P.R., Jr., and Foster, R.W., 1989, Future petroleum provinces in New Mexico--Discovering new reserves: New Mexico Bureau of Mines and Mineral Resources, 94 p. (17" x 22" folio), prepared for the New Mexico Research and Development Institute.
- Greenwood, E., Kottlowski, F.E., and Thompson, S., III, 1977, Petroleum potential and stratigraphy of Pedregosa Basin--Comparison with Permian and Orogrande Basins: American Association of Petroleum Geologists Bulletin, v. 61, p. 1448-1469.
- Grim, P.J., and Berry, G.W., 1979, Geothermal energy in the western United States, *in* Muffler, L.J.P., ed., Assessment of geothermal resources of the United States--1978: National Oceanic and Atmospheric Administration, National Geophysics Solar-Terrestrial Data Center and U.S. Geological Survey Circular 790, scale 1:2,500,000.

- Gurgel, K.D., ed., 1983, Energy resources map of Utah: Utah Geological and Mineral Survey Map 68, scale 1:500,000.
- Gustafson, V.O., 1981, Petroleum geology of the Devonian and Mississippian rocks of the Four Corners region, *in* Wiegand, D.L., ed., *Geology of the Paradox Basin: Rocky Mountain Association of Geologists Field Conference Guidebook*, p. 101-109.
- Guzman, A.B., 1981, Petroleum prospects in Mexico's Altar Desert: *Oil and Gas Journal*, v. 79, no. 34, August 24th, p. 110-126.
- Hamblin, W.K., 1965, Origin of "reverse drag" on the downthrown side of normal faults: *Geological Society of America Bulletin*, v. 76, p. 1145-1164.
- Harding, L.E., and Coney, P.J., 1985, The geology of the McCoy Mountains Formation, southeastern California and southwestern Arizona: *Geological Society of America Bulletin*, v. 96, p. 755-769.
- Harrison, G.C., 1976, Facies analysis of the Devonian of the Black Mesa Basin, Arizona: Flagstaff, Arizona, Northern Arizona University, unpublished M.S. thesis, 57 p.
- Hawley, J.W., and Lozinsky, R.P., 1992, Hydrogeologic framework of the Mesilla Basin in New Mexico and western Texas: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 323, 55 p.
- Hayes, P.T., 1970, Cretaceous paleogeography of southern Arizona and adjacent areas: U.S. Geological Survey Professional Paper 658-B, 42 p.
- Hayes, P.T., 1975, Cambrian and Ordovician rocks of southern Arizona and New Mexico and westernmost Texas: U.S. Geological Survey Professional Paper 873, 98 p.
- Hereford, Richard, 1977, Deposition of the Tapeats Sandstone (Cambrian) in central Arizona: *Geological Society of America Bulletin*, v. 88, p. 199-211.
- Heylmun, E.B., 1960, Kanab--The oil signs are there: *Oil and Gas Journal*, v. 58, no. 24, p. 168-171.
- Heylmun, E.B., 1981, Geology of promising Holbrook Basin: *Oil and Gas Journal*, v. 79, p. 117-118, 123.
- Hills, J.M., and Kottowski, F.E., coordinators, 1983, Correlation of stratigraphic units in North America--Southwest/southwest mid-continent correlation chart, Lindberg, F.A., ed.: American Association of Petroleum Geologists COSUNYA chart.

- Hintze, L.F., comp., 1980, Geologic map of Utah: Utah Geologic and Mineralogic Survey, scale 1:500,000.
- Hintze, L.F., 1988, Geologic history of Utah--A field guide to Utah's rocks: Brigham Young University Geology Studies Special Publication 7, 204 p.
- Huddle, J.W., and Dobrovolsky, E., 1952, Devonian and Mississippian rocks of central Arizona: U.S. Geological Survey Professional Paper 233-D, p. 67-112.
- Hunt, C.B., 1956, Cenozoic geology of the Colorado Plateau: U.S. Geological Survey Professional Paper 279, 99 p.
- Ijirigho, B.T., 1977, Pennsylvanian subsurface stratigraphy of the Black Mesa Basin and Four Corners area in northeastern Arizona: Tucson, Arizona, University of Arizona, unpublished M.S. thesis, 62 p.
- Jenney, J.P., and Stone, Claudia, eds., 1980, Studies in western Arizona: Arizona Geological Society Digest, V. XII, 322 p.
- Jenney, J.P., and Reynolds, S.J., eds., 1989, Geologic evolution of Arizona: Arizona Geological Society Digest 17, 866 p.
- Jordan, C.F., Jr., 1971, Lower Permian stratigraphy of southern New Mexico and West Texas: Houston, Texas, Rice University, unpublished Ph.D. dissertation, 136 p.
- Kashfi, M.S., 1983, Upper Devonian source and reservoir rocks in the Black Mesa Basin, northeastern Arizona: Oil and Gas Journal, v. 81, no. 50, December 12, p. 151-159.
- Kelley, V.C., and Clinton, N.J., 1960, Fracture systems and tectonic elements of the Colorado Plateau: University of New Mexico Publications in Geology no. 6, 104 p.
- Keith, S.B., and Wilt, J.C., 1986, Laramide orogeny in Arizona and adjacent regions--A strato-tectonic synthesis, *in* Beatty, B., and Wilkinson, P.A.K., eds., *Frontiers in geology and ore deposits of Arizona and the Southwest*: Arizona Geological Society Digest, V. XVI, p. 502-554.
- Kent, W.N., and Rawson, R.R., 1989, Depositional environments of the Mississippian Redwall Limestone in northeastern Arizona, *in* Fouch, T.D., and Magathan, E.R., eds., *Paleozoic paleogeography of west-central United States*: Society of Economic Paleontologists and Mineralogists, Symposium 1, p. 101-109.
- Kluth, C.F., 1986, Plate tectonics of the ancestral Rocky Mountains, *in* Peterson, J.A., ed., *Paleotectonics and sedimentation in the Rocky Mountain region, United States*: American Association of Petroleum Geologists Memoir 41, p. 353-369.

- Koons, E.D., 1945, Geology of the Uinkaret Plateau, northern Arizona: Geological Society of America Bulletin, v. 56, p. 151-180.
- Kottlowski, F.E., 1965, Sedimentary basins of south-central and south-western New Mexico: American Association of Petroleum Geologists Bulletin, v. 49, p. 2120-2139.
- Kottlowski, F.E., 1977, New Mexico geology and petroleum potential: Interstate Oil Compact Commission Committee Bulletin, v. 19, December, no. 2, p. 26-30.
- Kron, A., and Stix, J., 1982, Geothermal gradient map of the United States: National Oceanic and Atmospheric Administration, National Geophysics Data Center, prepared for Los Alamos National Laboratory, University of California, scale 1:2,500,000.
- Lessentine, R.H., 1965, Kaiparowits and Black Mesa Basin--Stratigraphic synthesis: American Association of Petroleum Geologists Bulletin, v. 49, p. 1997-2019.
- Lochman-Balk, C., 1972, Cambrian System, *in* Mallory, W.W., ed., Geologic atlas of the Rocky Mountain region: Rocky Mountain Association of Geologists, p. 60-75.
- Lovejoy, E.M.P., 1973, Major Early Cenozoic deformation along Hurricane Fault zone, Utah and Arizona: American Association of Petroleum Geologists Bulletin, v. 57, p. 510-519.
- Luedke, R.G., and Smith, R.L., 1978, Map showing distribution, composition, and age of Late Cenozoic volcanic centers in Arizona and New Mexico: U.S. Geological Survey Miscellaneous Investigations Series Map I-1091-A, scale 1:1,000,000.
- Lyth, A.L., Jr., 1971, Summary of possible future petroleum potential, region 3, Western Rocky Mountains, *in* Cram, I.H., ed., Future petroleum provinces of the United States--Their geology and potential: American Association of Petroleum Geologists Memoir 15, vol. 1, p. 406-412.
- Mack, G.H., Lawton, T.F., and Lucas, S.G., eds., 1988, Cretaceous and Laramide tectonic evolution of southwestern New Mexico: New Mexico Geological Society, 39th Field Conference Guidebook, 216 p.
- McKee, E.D., 1967, Arizona and western New Mexico, *in* Paleotectonic investigations of the Permian System in the United States: U.S. Geological Survey Professional Paper 515-J, p. 200-223.
- McKee, E.D., 1975, Arizona, *in* McKee, E.D., and Crosby, E.J., eds., Paleotectonic investigations of the Pennsylvanian System in the United States: U.S. Geological Survey Professional Paper 853-P, p. 293-309.

- McKee, E.D., 1979, Arizona, *in* Craig, L.C., and Conner, C.W., coordinators, Paleotectonic investigations of the Mississippian System in the United States, pt. 1: U.S. Geological Survey Professional Paper 1010-L, p. 198-207.
- McKee, E.D., 1982, The Supai Group of Grand Canyon: U.S. Geological Survey Professional Paper 1173, 504 p.
- McKenny, J.W., and Masters, J.A., 1968, Dineh-bi-Keyah Field, Apache County, Arizona: American Association of Petroleum Geologists Bulletin, v. 52, p. 2045-2057.
- Mallory, W.W., ed., 1972, Geologic atlas of the Rocky Mountain region: Rocky Mountain Association of Geologists, 331 p.
- Muehlberger, W.R., 1980, Texas lineament revisited, *in* Dickerson, P.W., and Hoffer, J.M., eds., Trans-Pecos region guidebook: New Mexico Geological Society 31st Field Conference, p. 113-121.
- Nations, J.D., Brennan, D.J., and Ybarra, R.A., 1989, Oil and gas in Arizona, *in* Jenney, J.P., and Reynolds, S.J., eds., Geologic evolution of Arizona: Arizona Geological Society Digest, V. 17, p. 795-816.
- Nations, J.D., Conway, C.M., and Swann, G.A., eds., 1986, Geology of central and northern Arizona: Geological Society of America Field Trip Guidebook for Rocky Mountain Section, Flagstaff, Arizona, Northern Arizona University, Geology Department, 176 p.
- Nations, J.D., Doss, A.K., and Ybarra, R.A., 1983a, Geologic setting of oil and gas exploration in Arizona, *in* Fassett, J.E., ed., Oil and gas fields of the Four Corners area, V. III: Four Corners Geological Society, p. 763-772.
- Nations, J.D., Doss, A.K., and Ybarra, R.A., 1983b, Stratigraphy and oil and gas production of Arizona, 1978-1983, *in* Fassett, J.E., ed., Oil and gas fields of the Four Corners area, V. III: Four Corners Geological Society, p. 832-843.
- Nations, J.D., and Eaton, J.G., eds., 1991, Stratigraphy, depositional environments, and sedimentary tectonics of the western margin, Cretaceous western Interior Seaway: Geological Society of America Special Paper 260.
- Nations, J.D., Landye, J.J., and Hevly, R.H., 1982, Location and chronology of Tertiary sedimentary deposits in Arizona--A review, *in* Ingersoll, R.V., and Woodburne, M.O., eds., Cenozoic nonmarine deposits of California and Arizona: Society of Economic Paleontologists and Mineralogists, Pacific Coast Section Symposium, p. 107-122.

- Nations, J.D., and Ranney, W.D.R., 1989, Geologic model of a Cenozoic basin in Arizona, *in* Tooker, E.W., ed., Arizona's industrial rock and mineral resources-- Workshop proceedings: U.S. Geological Survey Bulletin 1905, [in cooperation with Arizona Geological Survey], p. 44-53.
- Nations, J.D., and Stump, E., 1981, *Geology of Arizona*: Dubuque, Iowa, Kendall Hunt Publishing Company, 210 p.
- Nations, J.D., Wilt, J.C., and Hevly, R.H., 1985, Cenozoic paleogeography of Arizona, *in* Flores, R.M., and Kaplan, S.S., eds., *Cenozoic paleogeography of the west-central United States: Society of Economic Paleontologists and Mineralogists, Rocky Mountain Paleogeography Symposium 3*, p. 335-355.
- Oil and Gas Journal, 1979, Arizona area offers good potential for new finds: *Oil and Gas Journal*, v. 77, no. 44, p. 203-204.
- Olmsted, R.H., Loeltz, O.J., and Irelan, B., 1973, *Geohydrology of the Yuma area, Arizona and California*: U.S. Geological Survey Professional Paper 486-H, 227 p.
- Oppenheimer, J.M., and Sumner, J.S., 1980, *Depth-to-bedrock map, Basin and Range Province, Arizona*: Tucson, Arizona, University of Arizona, Department of Geosciences, scale 1:1,000,000.
- O'Sullivan, R.B., 1969, Crude oil, natural gas, and other associated gases, *in* *Mineral and water resources of Arizona*: Arizona Bureau of Mines Bulletin 180, p. 70-91.
- Peirce, H.W., 1976a, Tectonic significance of Basin and Range thick evaporite deposits: *Arizona Geological Society Digest V. X*, p. 325-339.
- Peirce, H.W., 1976b, Elements of Paleozoic tectonism in Arizona: *Arizona Geological Society Digest V. X*, p. 37-57.
- Peirce, H.W., 1979, The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States--Arizona: U.S. Geological Survey Professional Paper 1110-Z, 20 p.
- Peirce, H.W., 1981, Major Arizona salt deposits: *Arizona Bureau of Geology and Mineral Technology Fieldnotes*, v. 11, no. 4, p. 1-4.
- Peirce, H.W., 1982, The search for petroleum in Arizona: *Arizona Bureau of Geology and Mineral Technology, Fieldnotes*, v. 12, no. 2, p. 1-5.
- Peirce, H.W., and Gerrard, T.A., 1966, Evaporites of the Permian Holbrook Basin, Arizona, *in* Rau, J.L., ed., *Second symposium on salt*: Northern Ohio Geological Society, v. 1, p. 1-10.

- Peirce, H.W., Jones, Nile, and Rogers, Ralph, 1977, A survey of uranium favorability of Paleozoic rocks in the Mogollon Rim and Slope region east-central Arizona: Arizona Bureau of Geology and Mineral Technology Circular 19, 70 p.
- Peirce, H.W., Keith, S.B., and Wilt, J.C., 1970 (reprinted 1975), Coal, oil, natural gas, helium, and uranium in Arizona: Arizona Bureau of Mines Bulletin 182, 289 p. and 19 plates.
- Peirce, H.W., and Scurlock, J.R., 1972, Arizona well information: Arizona Bureau of Mines Bulletin 185, 195 p.
- Peterson, J.A., ed., 1986, Paleotectonics and sedimentation in the Rocky Mountain region, United States: American Association of Petroleum Geologists Memoir 41, 693 p.
- Plauchut, Bernard, Tardy, Marc, Blanchet, Rene, Zimmermann, Matthis, Cabezas, Pascal, Rosaz, Thierry, and Sosson, Marc, 1989, Etude Geologique des Cordilleres Nord-Americaines; Les lineaments du Texas et Caltam entre Cordilleres Americanines et Sierras Madres Mexicaines--Nature, origine et evolution structurale; Les montagnes rocheuses meridionales au centre-ouest du Nouveau-Mexique; Le passage des Cordilleres Nord-Americaines aux Sierras Madres Mexicaines le long du Texas lineament geologie du SW du Nouveau Mexique; Tectoniques superposees dans le sud-est de l'Arizona: New Mexico Bureau of Mines and Mineral Resources Open-File Report 385, 76 p.
- Poole, F.G., and 8 others, 1967, Devonian of the Southwestern United States, *in* Oswald, D.H., ed., International Symposium on the Devonian System: Society of Petroleum Geologists, Proceedings, v. 1, September 1967, Calgary, Canada, p. 879-912.
- Pray, L.C., 1977, Stratigraphic and structural features of the Sacramento Mountain escarpment, New Mexico, *in* Butler, J.H., ed., Geology of the Sacramento Mountains, Otero County, New Mexico: West Texas Geological Society Guidebook, Publication no. 1977-68, p. 73-89.
- Pye, W.D., 1967, Arizona--A new frontier to explore: Oil and Gas Journal, v. 65, May 8th, p. 168-172.
- Pyron, A.J., and Gray, J.H., 1985, Don't ignore the Orogrande; its potential is promising: World Oil, May, p. 59-67.
- Rauzi, S.L., 1990, Distribution of Proterozoic hydrocarbon source rock in northern Arizona and southern Utah: Phoenix, Arizona, Arizona Oil and Gas Conservation Commission Special Publication 5, 38 p.; map scale 1:500,000.

- Rauzi, S.L., 1992, Exploration well tests salt near Phoenix [Arizona], *in* Arizona Geology: Arizona Geological Survey, v. 22, no. 4, p. 910.
- Rauzi, S.L., 1993, Oil and natural gas occurrence in Arizona: Arizona Geological Survey Publication OG-35 (OGCC Chart C-1).
- Rauzi, S.L., 1994, Implications of live oil shows in eastern Arizona geothermal test: Arizona Geological Survey Open-File Report 94-1, 16 p; 12 maps, scale 1:500,000.
- Reif, D.M., and Robinson, J.P., 1981, Geophysical, geochemical, and petrographic data and regional correlation from the Arizona State A-1 well, Pinal County, Arizona: Arizona Geological Society Digest, V. XIII, p. 99-109.
- Reiter, M., Mansure, A.J., and Shearer, C., 1979, Geothermal characteristics of the Colorado Plateau: Tectonophysics, v. 61, p. 183-195.
- Reynolds, M.W., Palacas, J.G., and Elston, D.P., 1988, Potential petroleum source rocks in the Late Proterozoic Chuar Group, Grand Canyon, Arizona, *in* Carter, L.M.H., ed., USGS research on energy resources--1988, Program and abstracts: U.S. Geological Survey Circular 1025, p. 49-50.
- Reynolds, S.J., 1988, Geologic map of Arizona: Arizona Geological Survey Map 26, scale 1:1,000,000.
- Reynolds, S.J., Roddy, M.S., and Welty, J.W., 1986, Map of Paleozoic rocks and conodont alteration indices in Arizona: Arizona Bureau of Geology and Mineral Technology Open-File Report 86-1, scale 1:1,000,000.
- Reynolds, S.J., and Trapp, R.A., 1986, Computerized databases and database access programs of the Arizona Geological Survey: Arizona Geological Survey Open-File Report 86-7, 69 p.
- Reynolds, S.J., Welty, J.W., and Spencer, J.E., 1986, Volcanic history of Arizona: Arizona Bureau of Geology and Mineral Technology Field Notes, v. 16, no. 2, p. 1-5.
- Rieker, R.E., ed., 1979, Rio Grande Rift--Tectonics and magmatism: Washington, D.C., American Geophysical Union, 438 p.
- Robertson, J.M., and Broadhead, R.F., project directors, 1993, Atlas of major Rocky Mountain gas reservoirs: New Mexico Bureau of Mines and Mineral Resources (published for a consortium of agencies including Gas Research Institute, U.S. Department of Energy, Colorado Geological Survey, Utah Geological Survey, and Geological Survey of Wyoming), 206 p. (oversize).

- Ross, C.A., and Ross, J.R.P., 1986, Paleozoic paleotectonics and sedimentation in Arizona and New Mexico, *in* Peterson, J.A., ed., Paleotectonics and sedimentation in the Rocky Mountain region, United States: American Association of Petroleum Geologists Memoir 41, p. 653-668.
- Roybal, G.H., 1981, Facies development in a Lower Cretaceous coral-rudist patch reef (Mural Limestone, southeast Arizona): Rocky Mountain Association of Geologists, *The Mountain Geologist*, v. 18, no. 3, p. 46-56.
- Ryder, R.T., 1983a, Petroleum potential of wilderness lands in Arizona, *in* Miller, B.M., ed., Petroleum potential of wilderness lands in the western United States: U.S. Geological Survey Circular 902-C, p. 1-22.
- Ryder, R.T., 1983b, Petroleum potential of wilderness lands in New Mexico, *in* Miller, B.M., ed., Petroleum potential of wilderness lands in the Western United States: U.S. Geological Survey Circular 902-I, p. 1-33.
- Schmidt, Nancy, Reynolds, S.J., and Horstman, K.C., 1991, AZGEOBIB--A preliminary list of references on the geology of Arizona: Arizona Geological Survey Open-File Report 91-4, 302 p.
- Scurlock, J.R., 1971, Geologic structure map of northern Arizona: Phoenix, Arizona, Arizona Oil and Gas Conservation Commission, scale 1:375,000.
- Seager, W.R., 1983, Laramide wrench faults, basement-cored uplifts, and complementary basins in southern New Mexico: New Mexico Bureau of Mines and Mineral Resources, *New Mexico Geology*, v. 5, no. 4, p. 69-76.
- Seager, W.R., Clemons, R.E., and Callender, J.F., eds., 1975, Guidebook of the Las Cruces country: New Mexico Geological Society, 26th Field Conference, 345 p.
- Seager, W.R., and Morgan, P., 1979, Rio Grande Rift in southern New Mexico, West Texas, and northern Chihuahua, *in* Riecker, R.E., ed., Rio Grande Rift--Tectonism and magmatism: Washington, D.C., American Geophysical Union, p. 87-106.
- Shride, A.F., 1967, Younger Precambrian geology in southern Arizona: U.S. Geological Survey Professional Paper 566, 89 p.
- Smith, R.B., and Eaton, G.P., eds., 1978, Cenozoic tectonics and regional geophysics of the western Cordillera: Geological Society of America Memoir 152, 388 p.
- Stewart, W.J., ed., 1969, Delaware Basin exploration guidebook: West Texas Geological Society Publication no. 68-55a, 170 p.

- Stone, Claudia, and Jenney, J.P., eds., 1981: Arizona Geological Society Digest, V. XIII, 193 p.
- Stone, Claudia, and Witcher, J.C., 1982, Geothermal energy in Arizona--Final contract report: Arizona Bureau of Geology and Mineral Technology for the U.S. Department of Energy, Contract No. DE-FC07-79ID12009, 398 p.
- Sumner, J.S., 1975, Bouguer gravity map of Arizona: Tucson, Arizona, University of Arizona Laboratory of Geophysics, scale 1:1,000,000.
- Sumner, J.S., 1985, Crustal geology of Arizona as interpreted from magnetic, gravity, and geologic data, *in* Hinze, W.J., ed., The utility of regional gravity and magnetic anomaly maps: Society of Exploration Geophysicists, p. 164-180.
- Sylvester, A.G., comp., 1984, Wrench fault tectonics: American Association of Petroleum Geologists Reprint Series no. 23, 374 p.
- Teichert, C., 1965, Devonian rocks and paleogeography of central Arizona: U.S. Geological Survey Professional Paper 464, 181 p.
- Terra Graphics, 1977, Oil and gas production map of the western United States: Denver, Colorado, Terra Graphics, scale approximately 1:3,250,000.
- Thompson, G.A., and Zoback, M.L., 1979, Regional geophysics of the Colorado Plateau: Tectonophysics, v. 61, p. 149-181.
- Thompson, Sam, III, 1981, Analyses of petroleum source and reservoir rocks in southwestern New Mexico--Petroleum source rocks in exploration wells drilled to Paleozoic or Mesozoic units, Hidalgo and Grant Counties, New Mexico: New Mexico Bureau of Mines and Mineral Resources Open-File Report OF-153, 120 p.
- Thompson, Sam, III, 1982, Oil and gas exploration wells in southwestern New Mexico, *in* Powers, R.B., ed., Geologic studies of the Cordilleran thrust belt, v. II: Rocky Mountain Association of Geologists, p. 521-535.
- Thompson, Sam, III, and Bierberman, R.A., 1975, Oil and gas exploration wells in Do-a Ana County, New Mexico, *in* Seager, W.R., Clemons, R.E., and Callender, J.F., eds., Guidebook of the Las Cruces Country: New Mexico Geological Society, 26th Field Conference, p. 171-174.
- Thompson, Sam, III, and Jacka, A.D., 1981, Pennsylvanian stratigraphy, petrography, and petroleum geology of Big Hatchet Peak section, Hidalgo County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Circular 176, 129 p.

- Titley, S.R., ed., 1968, *Southern Arizona Guidebook III: Arizona Geological Society* (in cooperation with the Geological Society of America), 354 p.
- Tooker, E.W., comp.-ed., 1989, *Arizona's industrial rock and mineral resources-- Workshop proceedings: U.S. Geological Survey Bulletin 1905* (with the Arizona Geological Survey), 93 p.
- Tovar R, J.C., 1981, *Provincias con posibilidades petroleras en el Distrito de Chihuahua: Boletin de la Asociacion Mexicana de Geologos Petroleros*, v. 33, no. 1, p. 25-51.
- Tucker, M.L., 1983, *Geochemical correlation of crude oil and source rocks in Black Mesa Basin, northeastern Arizona: Flagstaff, Arizona, Northern Arizona University*, unpublished M.S. thesis, 80 p.
- Turner, D.S., 1968, *Natural gas in Black Mesa Basin, northeastern Arizona*, in Beebe, W., ed., *Natural gases of North America: American Association of Petroleum Geologists Memoir 9*, v. 2, p. 1357-1370.
- Wardlaw, B.R., and Harris, A.G., 1984, *Conodont-based thermal maturation of Paleozoic rocks in Arizona: American Association of Petroleum Geologists Bulletin*, v. 69, p. 1101-1106.
- Weiss, Emanuel, 1991, *Regional ground-water flow in upper and middle Paleozoic rocks in southeastern Utah and adjacent parts of Arizona, Colorado, and New Mexico: U.S. Geological Survey Water-Resources Investigations Report 90-4079*, 57 p.
- Wengerd, S.A., 1970, *Petroleum prospects in southwesternmost New Mexico*, in Woodward, L.A., ed., *Guidebook of the Tyrone--Big Hatchet Mountains Florida Mountains region: New Mexico Geological Society, 21st Field Conference*, p. 91-104.
- Wilson, E.D., 1962, *A r̄sum̄ of the geology of Arizona: Arizona Bureau of Mines Bulletin 171*, 140 p.
- Wilson, E.D., and Moore, R.T., 1962, *Geologic cross-sections of Arizona 1-8: Arizona Bureau of Mines*, horizontal scale 1 inch equals 3 miles, vertical scale 1 inch equals 500 feet.
- Wilson, E.D., Moore, R.T., and Cooper, J.R., 1969, *Geologic map of Arizona: Arizona Bureau of Mines and U.S. Geological Survey*, scale 1:500,000.
- Wilson, J.L., 1989, *Lower and Middle Pennsylvanian strata in the Orogrande and Pedregosa Basins, New Mexico: New Mexico Bureau of Geology and Mineral Resources Bulletin 124*, 16 p.

- Wilt, J.C., and Jenney, J.P., eds., 1976, Tectonic digest: Arizona Geological Society digest, V. X, 430 p.; 4 plates.
- Woodward-Clyde Consultants, 1983, Overview of the regional geology of the Paradox Basin study region, *for* the U.S. Department of Energy, Office of Nuclear Waste Isolation, ONWI-92, distribution category UC-70: San Francisco, California, 433 p.
- Woodward, L.A., ed., 1970, Guidebook of the Tyrone--Big Hatchet Mountains Florida Mountains region: New Mexico Geological Society, 21st Field Conference, 162 p.
- Woodward, L.A., 1973, Structural framework and tectonic evolution of the Four Corners region of the Colorado Plateau, *in* James, H.L., ed., Guidebook of Monument Valley and vicinity Arizona and Utah: New Mexico Geological Society, 24th Field Conference, p. 94-98.
- Woodward, L.A., comp., 1984, Tectonic map of the Rocky Mountain region of the United States, *in* Sloss, L.L., ed., Sedimentary cover--North American craton: Geological Society of America Decade of North American Geology Project, v. D-2, plate 2, scale 1:2,500,000.
- Woodward, L.A., and Duchene, H.R., 1981, Overthrust belt of southwestern New Mexico: Comparison with Wyoming-Utah overthrust belt: American Association of Petroleum Geologists Bulletin, v. 65, p. 722-729.
- Woodward, L.A., and Grant, P.R., 1986, Central-western New Mexico--An exploration frontier for oil and gas, *in* Clemons, R.E., King, W.E., and Mack, G.H., eds., Truth or Consequences region: New Mexico Geological Society Guidebook, 37th Field Conference, p. 307-314.
- Woodward, L.A., and Northrop, S.A., eds., 1976, Tectonics and mineral resources of Southwestern North America: New Mexico Geological Society Special Publication no. 6, 218 p.
- Wrucke, C.T., and Armstrong, A.K., 1987, Paleozoic stratigraphic section in Dry Canyon, Whetstone Mountains, Cochise County, Arizona, *in* Hill, M.L., ed., Centennial field guide volume 1: Geological Society of America, Cordilleran Section, p. 29-34.

- Wrucke, C.T., Otton, J.K., and Desborough, G.A., 1986, Summary and origin of the mineral commodities in the Middle Proterozoic Apache Group in central Arizona, *in* Geatty, B., and Wilkinson, P.A.K., eds., *Frontiers in geology and ore deposits of Arizona and the Southwest: Arizona Geological Society Digest*, V. XVI, p. 12-17.
- Young, R.A., 1979, Laramide deformation, erosion, and plutonism along the southwestern margin of the Colorado Plateau: *Tectonophysics*, v. 61, p. 25-47.
- Zietz, I., comp., 1982, Composite magnetic anomaly map of the United States: U.S. Geological Survey and Society of Exploration Geophysicists, scale 1:2,500,000.